

IDENTITIES OF SYMMETRY FOR HIGHER-ORDER q -EULER POLYNOMIALS

DAE SAN KIM AND TAEKYUN KIM

ABSTRACT. In this paper, we derive basic identities of symmetry in two variables related to higher-order q -Euler polynomials and q -analogue of higher-order alternating power sums. The derivation of identities are based on the multivariate p -adic fermionic integral expression of the generating function for the higher-order q -Euler polynomials.

1. INTRODUCTION

Let p be a fixed prime number. Throughout this paper, $\mathbb{Z}_p$, $\mathbb{Q}_p$ and $\mathbb{C}_p$ will, respectively, denote the ring of p -adic rational integers, the field of p -adic rational numbers and the completion of algebraic closure of $\mathbb{Q}_p$.

Let v_p be the normalized exponential valuation of $\mathbb{C}_p$ with $|p|_p = p^{-v_p(p)} = \frac{1}{p}$. When one talks about q -extension, q is variously considered as an indeterminate, a complex number $q \in \mathbb{C}$ or p -adic number $q \in \mathbb{C}_p$. If $q \in \mathbb{C}$, one usually assumes $|q| < 1$; if $q \in \mathbb{C}_p$, one usually assumes $|1 - q|_p < 1$. The q -number of x is defined by $[x]_q = \frac{1-q^x}{1-q}$. Note that $\lim_{q \rightarrow 1} [x]_q = x$.

For $r \in \mathbb{N}$, the Euler polynomials of order r are defined by the generating function to be

$$(1) \quad \left(\frac{2}{e^t + 1} \right)^r e^{xt} = \left(\frac{2}{e^t + 1} \right) \times \cdots \times \left(\frac{2}{e^t + 1} \right) e^{xt} = \sum_{n=0}^{\infty} E_n^{(r)}(x) \frac{t^n}{n!}.$$

When $x = 0$, $E_n = E_n(0)$ are called the Euler numbers of order r (see [?, ?]).

From (1), we note that

$$(2) \quad E_n^{(r)}(x) = \sum_{l=0}^n \binom{n}{l} E_l^{(r)} x^{n-l},$$

(see [?, ?]).

In [?], Kim considered the q -extension of (1) which is given by

$$(3) \quad 2^r \sum_{m=0}^{\infty} (-1)^m \binom{m+r-1}{m} e^{[m+x]_q t} = \sum_{n=0}^{\infty} E_{n,q}^{(r)}(x) \frac{t^n}{n!}.$$

Thus, by (3), we get

$$(4) \quad \sum_{n=0}^{\infty} E_{n,q}^{(r)}(x) \frac{t^n}{n!} = 2^r \sum_{m_1, \dots, m_r=0}^{\infty} (-1)^{m_1+\dots+m_r} e^{[m_1+\dots+m_r+x]_q t},$$

where $E_{n,q}^{(r)}(x)$ are called the q -Euler polynomials of order $r \in \mathbb{N}$.

When $x = 0$, $E_{n,q}^{(r)} = E_{n,q}^{(r)}(0)$ are called the q -Euler number of order r .
From (4), we note that

$$(5) \quad E_{n,q}^{(r)}(x) = \sum_{l=0}^n \binom{n}{l} q^{lx} E_{l,q}^{(r)} [x]_q^{n-l}.$$

Let $\mathcal{C}(\mathbb{Z}_p)$ be the space of continuous functions on $\mathbb{Z}_p$. For $f \in \mathcal{C}(\mathbb{Z}_p)$, the fermionic p -adic integral on $\mathbb{Z}_p$ is defined by Kim as follows :

$$(6) \quad I_{-1}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x) = \lim_{N \rightarrow \infty} \sum_{x=0}^{p^N-1} f(x) (-1)^x,$$

(see [?, ?, ?]).

By (6), we easily geet

$$(7) \quad \int_{\mathbb{Z}_p} f(x+n) d\mu_{-1}(x) + (-1)^{n-1} \int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x) = 2 \sum_{l=0}^{n-1} (-1)^{n-1-l} f(l),$$

where $n \in \mathbb{N}$ (see [?, ?, ?, ?, ?, ?]).

Thus, from (7), we have

$$(8) \quad \int_{\mathbb{Z}_p} e^{(x+y)t} d\mu_{-1}(y) = \frac{2}{e^t + 1} e^{xt} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}.$$

By (8), we easily get

$$(9) \quad \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} e^{(x+y_1+\cdots+y_r)t} d\mu_{-1}(y_1) \cdots d\mu_{-1}(y_r) = \left(\frac{2}{e^t + 1} \right)^r e^{xt} \\ = \sum_{n=0}^{\infty} E_n^{(r)}(x) \frac{t^n}{n!}.$$

In the next section, we consider the q -analogue of (9). In [?], the thirty one basic identities of symmetry in three variables related to higher-order Euler polynomials and alternating power sums are derived from (9).

In [?], Kim gave some interesting relations of symmetry between the alternating power sum polynomials and Euler polynomials and he suggested an open question as to finding the q -extension of symetry p -adic invariant integral on $\mathbb{Z}_p$ for q -Euler polynomials.

Recently, several authors have studied the identity of symmetry and q -extensions of Euler polynomials which are derived from the p -adic fermionic integrals on $\mathbb{Z}_p$ (see [1-13]).

In this paper, we investigate several further interesting properties of symmetry for the multivariate p -adic fermionic integrals on q -polynomials. From our investigation, we derive some relations of symmetry between the higher-order alternating power sum q -polynomials and the higher-order q -Euler polynomials.

2. IDENTITIES OF SYMMETRY FOR THE HIGHER-ORDER q -EULER POLYNOMIALS

From (3), (4), (6) and (7), we note that

$$\begin{aligned}
 (10) \quad & \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} e^{[x+y_1+\cdots+y_r]_q t} d\mu_{-1}(y_1) \cdots d\mu_{-1}(y_r) \\
 &= 2^r \sum_{m_1, \dots, m_r=0}^{\infty} (-1)^{m_1+\cdots+m_r} e^{[m_1+\cdots+m+x]_q t} \\
 &= \sum_{n=0}^{\infty} E_{n,q}^{(r)}(x) \frac{t^n}{n!}.
 \end{aligned}$$

Thus, by (10), we get

$$\begin{aligned}
 (11) \quad & \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [x+y_1+\cdots+y_r]_q^n d\mu_{-1}(y) \\
 &= E_{n,q}^{(r)}(x) \\
 &= \frac{1}{(1-q)^n} \sum_{l=1}^n \binom{n}{l} (-1)^l \left(\frac{2}{1+q^l} \right)^r \\
 &= 2^r \sum_{m_1, \dots, m_r=0}^{\infty} (-1)^{m_1+\cdots+m_r} [m_1+\cdots+m_r]_q^n.
 \end{aligned}$$

Let $w_1, w_2 \in \mathbb{N}$ with $w_1 \equiv 1 \pmod{2}$ and $w_2 \equiv 1 \pmod{2}$. Then we observe that

$$\begin{aligned}
 (12) \quad & \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} e^{[w_1]_q [w_2 x + \frac{w_2}{w_1} (j_1 + \cdots + j_r) + y_1 + \cdots + y_r]_q} e^{w_1 t} d\mu_{-1}(y_1) \cdots d\mu_{-1}(y_r) \\
 &= \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} e^{[w_1 w_2 x + w_2 (j_1 + \cdots + j_r) + w_1 (y_1 + \cdots + y_r)]_q} e^{t} d\mu_{-1}(y_1) \cdots d\mu_{-1}(y_r) \\
 &= \lim_{N \rightarrow \infty} \sum_{y_1, \dots, y_r=0}^{p^N-1} e^{[w_1 w_2 x + w_2 (j_1 + \cdots + j_r) + w_1 (y_1 + \cdots + y_r)]_q} (-1)^{y_1 + \cdots + y_r} \\
 &= \lim_{N \rightarrow \infty} \sum_{y_1, \dots, y_r=0}^{w_2 p^N-1} e^{[w_1 w_2 x + w_2 \sum_{l=1}^r j_l + w_1 \sum_{l=1}^r y_l]_q} (-1)^{\sum_{l=1}^r y_l} \\
 &= \lim_{N \rightarrow \infty} \sum_{i_1, \dots, i_r=0}^{w_2-1} \sum_{y_1, \dots, y_r=0}^{p^N-1} e^{[w_1 w_2 x + w_2 \sum_{l=1}^r j_l + w_1 \sum_{l=1}^r (i_l + w_2 y_l)]_q} (-1)^{\sum_{l=1}^r (i_l + w_2 y_l)}.
 \end{aligned}$$

From (12), we have

$$\begin{aligned}
(13) \quad & \sum_{j_1, \dots, j_r=0}^{w_1-1} (-1)^{j_1+\dots+j_r} \\
& \times \int_{\mathbb{Z}_p} \dots \int_{\mathbb{Z}_p} e^{[w_1]_q \left[w_2 x + \frac{w_2}{w_1} (j_1 + \dots + j_r) + y_1 + \dots + y_r \right]_q w_1 t} d\mu_{-1}(y_1) \dots d\mu_{-1}(y_r) \\
& = \lim_{N \rightarrow \infty} \left(\sum_{j_1, \dots, j_r=0}^{w_1-1} \sum_{i_1, \dots, i_r=0}^{w_2-1} \sum_{y_1, \dots, y_r=0}^{p^N-1} (-1)^{\sum_{l=1}^r (i_l + j_l + y_l)} \right. \\
& \quad \left. \times e^{[w_1 w_2 (x + \sum_{l=1}^r y_l) + w_2 \sum_{l=1}^r j_l + w_1 \sum_{l=1}^r i_l]_q t} \right).
\end{aligned}$$

By the same method as (13), we get

$$\begin{aligned}
(14) \quad & \sum_{j_1, \dots, j_r=0}^{w_2-1} (-1)^{j_1+\dots+j_r} \\
& \times \int_{\mathbb{Z}_p} \dots \int_{\mathbb{Z}_p} e^{[w_2]_q \left[w_1 x + \frac{w_1}{w_2} (j_1 + \dots + j_r) + y_1 + \dots + y_r \right]_q w_2 t} d\mu_{-1}(y_1) \dots d\mu_{-1}(y_r) \\
& = \lim_{N \rightarrow \infty} \left(\sum_{j_1, \dots, j_r=0}^{w_2-1} \sum_{i_1, \dots, i_r=0}^{w_1-1} \sum_{y_1, \dots, y_r=0}^{p^N-1} (-1)^{\sum_{l=1}^r (i_l + j_l + y_l)} \right. \\
& \quad \left. \times e^{[w_1 w_2 (x + \sum_{l=1}^r y_l) + w_1 \sum_{l=1}^r j_l + w_2 \sum_{l=1}^r i_l]_q t} \right).
\end{aligned}$$

Therefore, by (13) and (14), we obtain the following theorem.

Theorem 1. For $w_1, w_2 \in \mathbb{N}$ with $w_1 \equiv 1 \pmod{2}$ and $w_2 \equiv 1 \pmod{2}$, we have

$$\begin{aligned}
& \sum_{j_1, \dots, j_r=0}^{w_1-1} (-1)^{j_1+\dots+j_r} \\
& \times \int_{\mathbb{Z}_p} \dots \int_{\mathbb{Z}_p} e^{[w_1]_q \left[w_2 x + \frac{w_2}{w_1} (j_1 + \dots + j_r) + y_1 + \dots + y_r \right]_q w_1 t} d\mu_{-1}(y_1) \dots d\mu_{-1}(y_r) \\
& = \sum_{j_1, \dots, j_r=0}^{w_2-1} (-1)^{j_1+\dots+j_r} \\
& \times \int_{\mathbb{Z}_p} \dots \int_{\mathbb{Z}_p} e^{[w_2]_q \left[w_1 x + \frac{w_1}{w_2} (j_1 + \dots + j_r) + y_1 + \dots + y_r \right]_q w_2 t} d\mu_{-1}(y_1) \dots d\mu_{-1}(y_r).
\end{aligned}$$

Corollary 2. For $n \geq 0$, and $w_1, w_2 \in \mathbb{N}$ with $w_1 \equiv 1 \pmod{2}$ and $w_2 \equiv 1 \pmod{2}$, we have

$$\begin{aligned} & [w_1]_q^n \sum_{j_1, \dots, j_r=0}^{w_1-1} (-1)^{\sum_{l=1}^r j_l} \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} \left\{ \left[w_2 x + \frac{w_2}{w_1} (j_1 + \cdots + j_r) + (y_1 + \cdots + y_r) \right]_{q^{w_1}}^n \right\} d\mu_{-1}(y_1) \cdots d\mu_{-1}(y_r) \\ &= [w_2]_q^n \sum_{j_1, \dots, j_r=0}^{w_2-1} (-1)^{\sum_{l=1}^r j_l} \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} \left\{ \left[w_1 x + \frac{w_1}{w_2} (j_1 + \cdots + j_r) + (y_1 + \cdots + y_r) \right]_{q^{w_2}}^n \right\} d\mu_{-1}(y_1) \cdots d\mu_{-1}(y_r). \end{aligned}$$

Therefore, by (11) and Corollary 2, we obtain the following theorem.

Theorem 3. For $n \geq 0$, and $w_1, w_2 \in \mathbb{N}$ with $w_1 \equiv 1 \pmod{2}$ and $w_2 \equiv 1 \pmod{2}$, we have

$$\begin{aligned} & [w_1]_q^n \sum_{j_1, \dots, j_r=0}^{w_1-1} (-1)^{\sum_{l=1}^r j_l} E_{n, q^{w_1}}^{(r)} \left(w_2 x + \frac{w_2}{w_1} (j_1 + \cdots + j_r) \right) \\ &= [w_2]_q^n \sum_{j_1, \dots, j_r=0}^{w_2-1} (-1)^{\sum_{l=1}^r j_l} E_{n, q^{w_2}}^{(r)} \left(w_1 x + \frac{w_1}{w_2} (j_1 + \cdots + j_r) \right). \end{aligned}$$

From (11), we can derive the following equation (15) :

$$\begin{aligned} (15) \quad & \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} \left[w_2 x + \frac{w_2}{w_1} (j_1 + \cdots + j_r) + (y_1 + \cdots + y_r) \right]_{q^{w_1}}^n d\mu_{-1}(y_1) \cdots d\mu_{-1}(y_r) \\ &= \sum_{i=0}^n \binom{n}{i} \left(\frac{[w_2]_q}{[w_1]_q} \right)^i [j_1 + \cdots + j_r]_{q^{w_2}}^i q^{w_2(n-i) \sum_{l=1}^r j_l} \\ & \quad \times \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} \left[w_2 x + \sum_{l=1}^r y_l \right]_q^{n-i} d\mu_{-1}(y_1) \cdots d\mu_{-1}(y_r) \\ &= \sum_{i=0}^n \binom{n}{i} \left(\frac{[w_2]_q}{[w_1]_q} \right)^i [j_1 + \cdots + j_r]_{q^{w_2}}^i q^{w_2(n-i) \sum_{l=1}^r j_l} E_{n-i, q^{w_1}}^{(r)}(w_2 x). \end{aligned}$$

By (15), we get

$$\begin{aligned}
(16) \quad & [w_1]_q^n \sum_{j_1, \dots, j_r=0}^{w_1-1} (-1)^{\sum_{l=1}^r j_l} \\
& \times \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} \left[w_2 x + \frac{w_2}{w_1} \sum_{l=1}^r j_l + \sum_{l=1}^r y_l \right]_{q^{w_1}}^n d\mu_{-1}(y_1) \cdots d\mu_{-1}(y_r) \\
& = \sum_{j_1, \dots, j_r=0}^{w_1-1} (-1)^{\sum_{l=1}^r j_l} \sum_{i=0}^n \binom{n}{i} [w_2]_q^i [w_1]_q^{n-i} \\
& \quad \times [j_1 + \cdots + j_r]_{q^{w_2}}^i q^{w_2(n-i)} \sum_{l=1}^r j_l E_{n-i, q^{w_1}}^{(r)}(w_2 x) \\
& = \sum_{i=0}^n \binom{n}{i} [w_1]_q^{n-i} [w_2]_q^i E_{n-i, q^{w_1}}^{(r)}(w_2 x) \\
& \quad \times \sum_{j_1, \dots, j_r=0}^{w_1-1} (-1)^{j_1 + \cdots + j_r} [j_1 + \cdots + j_r]_{q^{w_2}}^i q^{w_2(n-i)} \sum_{l=1}^r j_l \\
& = \sum_{i=0}^n \binom{n}{i} [w_1]_q^{n-i} [w_2]_q^i T_{n, i, q^{w_2}}^{(r)}(w_1) E_{n-i, q^{w_1}}^{(r)}(w_2 x),
\end{aligned}$$

where

$$(17) \quad T_{n, i, q}^{(r)}(w) = \sum_{j_1, \dots, j_r=0}^{w-1} (-1)^{j_1 + \cdots + j_r} q^{(n-i)(j_1 + \cdots + j_r)} [j_1 + \cdots + j_r]_q^i.$$

By the same method as (16), we see that

$$\begin{aligned}
(18) \quad & [w_2]_q^n \sum_{j_1, \dots, j_r=0}^{w_2-1} (-1)^{j_1 + \cdots + j_r} \\
& \times \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} \left[w_1 x + \frac{w_1}{w_2} (j_1 + \cdots + j_r) + (y_1 + \cdots + y_r) \right]_{q^{w_2}}^n d\mu_{-1}(y_1) \cdots d\mu_{-1}(y_r) \\
& = \sum_{i=0}^n \binom{n}{i} [w_1]_q^i [w_2]_q^{n-i} T_{n, i, q^{w_1}}^{(r)}(w_2) E_{n-i, q^{w_2}}^{(r)}(w_1 x).
\end{aligned}$$

Therefore, by Corollary (2), (16), (17) and (18), we obtain the following theorem.

Theorem 4. For $n \geq 0$, and $w_1, w_2 \in \mathbb{N}$ with $w_1 \equiv 1 \pmod{2}$ and $w_2 \equiv 1 \pmod{2}$, we have

$$\begin{aligned}
& \sum_{i=0}^n \binom{n}{i} [w_1]_q^{n-i} [w_2]_q^i T_{n, i, q^{w_2}}^{(r)}(w_1) E_{n-i, q^{w_1}}^{(r)}(w_2 x) \\
& = \sum_{i=0}^n \binom{n}{i} [w_1]_q^i [w_2]_q^{n-i} T_{n, i, q^{w_1}}^{(r)}(w_2) E_{n-i, q^{w_2}}^{(r)}(w_1 x),
\end{aligned}$$

where

$$T_{n, i, q}^{(r)}(w) = \sum_{j_1, \dots, j_r=0}^{w-1} (-1)^{j_1 + \cdots + j_r} q^{(n-i)(j_1 + \cdots + j_r)} [j_1 + \cdots + j_r]_q^i.$$

ACKNOWLEDGEMENTS. This work was supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government(MOE) (No.2012R1A1A2003786).

DEPARTMENT OF MATHEMATICS, SOGANG UNIVERSITY, SEOUL 121-742, REPUBLIC OF KOREA

E-mail address : `dskim@sogang.ac.kr`

DEPARTMENT OF MATHEMATICS, KWANGWOON UNIVERSITY, SEOUL 139-701, REPUBLIC OF KOREA

E-mail address : `tkkim@kw.ac.kr`